

*Ethnobotanical Leaflets* 12: 776-83. 2008.

## Effects of Some Heavy Metals on Chlorophyll Accumulation in *Barbula lambarenensis*

P.O. Fatoba and Udoh, Emem G.

Department of Plant Biology  
University of Ilorin  
Ilorin, Kwara State, Nigeria

Issued 04 October 2008

### ABSTRACT

Samples of moss (*Barbula lambarenensis*) with their substrates collected from Belewu Drive, Oke Odo, Ilorin were taken to the screen house of the University of Ilorin biological garden to monitor the effects and tolerance of this plant to different heavy metals. The moss samples were divided into eleven regimes, widely separated from one another to avoid contaminations. Ten regimes were differently irrigated with 1000ppm and 2000ppm of lead, copper, cadmium, iron and vanadium thrice a week. The eleventh treatment served as the control, and was irrigated with distilled water. It was found that these heavy metals caused some damage to the chloroplasts of this plant as the bright green colours changed light green, yellowish green or brown. The concentrations of the used heavy metals are phytotoxic. In light of this, *Barbula lambarenensis* can serve as a bio-indicator of heavy metals.

### INTRODUCTION

Many plants and animals have been used to assess environmental pollution (Kasanen and Venetvaara 1991). Of all, mosses and lichens have been found to be the best bio indicators (Murkherjee and Nuorteva, 1984). The ability of mosses in this regard could be attributed to their small-size, nakedness, habitat, poikilohydric nature and reduced sensitivity to the environment. Generally, bryophytes are well adapted to moist habitats while some are able to survive in areas with little or no rainfall (Carrington, 1997). Bryophytes, especially mosses are ubiquitous as they are equipped with some structural adaptive strategies that enable them to grow successfully where they occur. Their ability to grow on substrates or areas which are inhospitable to higher plants exclude them from intense competition and this gives them an added advantage to serve as good bio indicators. Gradzinski (1990) identified three types of bio indicators: true indicator (pollutants damage the plant proportionally to the concentration of the pollutants); accumulators (accumulate the potentially toxic pollutants); and scales of indicator species (pollutants lead to the disappearance or appearance of certain species).

Environmental pollution has become a matter that requires immediate attention as it affects human health and results in death. Urbanization, civilization and industrialization have worsened this ugly situation of the environment such that there is no more free air again. The pollutants are released from human activities such as agricultural practices (usage of pesticides, fungicides, fertilizers, etc), combustion, sewage and waste disposal, vehicular wear and tear, mining and smelting processes, tetra oxosulphate VI and trioxo nitrate V acid plants. All these produce poisonous pollutants such as SO<sub>2</sub>, NO<sub>x</sub>, heavy metals, SO<sub>3</sub> that affect both plants and animals. Critical about these pollutants

is the ill health associated with the lungs, heart and liver leading to death.

Bates (2000, 2002) reported that mosses are the least sensitive to the major pollutants of the environment. Kakulu (1993) reported that *Polytrichum junipercum* found on the bark of *Azadrachta indica* had been used to assess heavy metal pollution. He confirmed that the concentrations of pollutants in the tree bark correlate with those of the environment. Charaborty *et al* (2006) worked on the use of mosses as bio-monitor of trace element. It has been found that bryophytes growing directly on an alkaline masonry showed enhanced tolerance to SO<sub>2</sub> (Gilbert, 1970). This ability was assumed to be due to CaCO<sub>3</sub> concentration of the substrate which was believed to have neutralised the deposition of acid in the soil and pushed the toxic equilibrium from SO<sub>2</sub> dissolving in surface water solution towards sulphite, the least harmful of the three ion forms of SO<sub>2</sub> ; SO<sub>3</sub><sup>2-</sup>, HSO<sub>3</sub><sup>-</sup>, H<sub>2</sub>SO<sub>3</sub> in the soil. The toxicity of SO<sub>2</sub> results from the fact that the gas forms highly reactive species on dissolving in water forming SO<sub>3</sub><sup>2-</sup>, HSO<sub>3</sub><sup>-</sup>, H<sub>2</sub>SO<sub>3</sub>, the most toxic being HSO<sub>3</sub><sup>-</sup>, H<sub>2</sub>SO<sub>3</sub> (Richardson, 1981).

Most of the pollutants are silent killers, the need to identify bio -indicators in the environment is required as this would reveal the pollution status of man's environment from time to time. This would equally help in identifying the prevailing danger and thus enable man to take necessary step by checkmating the release of such pollutant and advise the government or people on the necessary steps to take so as to enjoy longevity and good health. This study was designed to identify the effects of some heavy metals on chlorophyll accumulation in *Barbula lambarenensis*.

## MATERIALS AND METHODS

*Barbula lambarenensis* (Pottiaceae) is a bright green, cushion like acrocarpous moss that grows on sandcrete materials under tree canopy or in cool environment supplied with adequate moisture. Although it reproduces sexually but rely mostly on asexual propagules, gemmae for its establishment and expansion (Fatoba 1998). This plant was chosen for this study because of its occurrence in the nooks and corners of the environment *B. lambarenensis* with its sandcrete substrates were collected along Yinka Belewu Drive, Oke-Odo, Tanke, Ilorin Kwara State, Nigeria These plant collections were taken to the screen house in the University of Ilorin Biological Garden where they were divided into eleven regimes with wide spaces left in between regimes to avoid contaminations and mix-up.

Five different salts: Pb (NO<sub>3</sub>)<sub>2</sub>, CdSO<sub>4</sub>, CuSO<sub>4</sub>, Fe<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub> and Va were used for the experiment to furnish five different heavy metals, Required amounts of each salt was weighed and dissolved in a 1litre of distilled water to give 5 and 10 mg/l of each of the metals (Table 1).

**Table 1. Preparation of working solutions.**

concentration Quantity of salts dissolved in 1 litre of distilled					
	Pb(NO <sub>3</sub> ) <sub>2</sub>	CdSO <sub>4</sub>	CUSO <sub>4</sub>	Fe <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub>	Va
5mg/l	0.00799	0.09270	0.01255	0.01788	0.00500
10mg/l	0.01599	0.01854	0.02510	0.03576	0.01000
Control (water)	-	-	-	-	-

Samples were collected from each regime weekly for chlorophyll determination and fortnightly for microscopic observations. Plant materials collected were extracted separately with 80% acetone, centrifuged and the optical densities read off at 645 and 660 wavelengths of the Spectrophotometer model Campspec M105. The chlorophylls a and b contents of each

sample determined with Arnon (1949) formula and total chlorophyll with Strain and Svec (1966) formula. The experiment lasted for three months and the data generated were subjected to Student t – paired test.

## OBSERVATIONS AND RESULTS

All the plants irrigated with heavy metal solutions have their bright green colours changed to light green, yellow or brown while the control remained green throughout the period of study. This shows that the heavy metals affected the chloroplast negatively. The change in colouration started at the 4<sup>th</sup> week of experiment in regimes irrigated with 5mg/l Cd, Fe and Pb, 6<sup>th</sup> week in regimes irrigated with Va and 8<sup>th</sup> week in samples irrigated with Cu.. The change was noticed earlier (4<sup>th</sup> week) in the regimes irrigated with 10mg/l of all the heavy metal solutions except in the regime irrigated with Va that showed in 6<sup>th</sup> week. The trend of this result showed that Cd, Feiii and Pb are more toxic/ and deleterious to the chloroplast than Va and Cu. The cells and nuclei of the control experiment remained intact throughout the period of study but the cell started collapsing and becoming distorted during the 4<sup>th</sup> week for Va treatment, 6<sup>th</sup> week for Cu, Cd, Fe and Pb treatments. The same trend

was observed for the nature of the nuclei except for the Cu treated samples. However, the effects started earlier (4<sup>th</sup> week) in regimes treated with 10mgll working solutions. In light of this, Va is more toxic than other heavy metals on the foliar cells and nuclei of the plant.

Generally, the trend of toxicity was Va> Cd Pb, Fe iii >Va> Cd> Cu as the foliar colours changed brown, light yellow and pale green respectively.

Table 2 shows the weekly effect of the different heavy metals on chl;orophyll a content of the treated *Barbula lambarenensis*. The treated samples had less chlorophyll a than the control (Table 2). Generally, the chlorophyll a contents of samples irrigated with 5mg/l of heavy metal solution were greater than those of 10mg/l. Both the control and treated followed same trend in respect of the chlorophyll a content. The chlorophyll a increased from the 1<sup>st</sup> week of experiment to the 5<sup>th</sup> week then gradually decreased to the end of the experiment. Chlorophyll a in the treated samples (5 and 10mgll) were less than that of the control after the 4<sup>th</sup> week of the experiment with some being greater than the control during the 1<sup>st</sup> & 2<sup>nd</sup> week (Cd, Fe, Pb) while Cu & Va were less for 5mg/l However, the pattern of the relationships of the treated with the control were not clearly defined for 2<sup>nd</sup> to 4<sup>th</sup> week (Table 2). Moreover, only the samples treated with 10mgll Fe and Pb had greater chlorophyll a than the control (Table 2). Table 2 further revealed the high chlorophyll a content of samples irrigated with both 5mgll and 10mgll of Cu f rom 2<sup>nd</sup> week to 4<sup>th</sup> week.

The mean chlorophyll a contents of the samples irrigated with 5mgll & 10mg/l of Cu and Pb were statistically the same with that of the control. This shows that these two heavy metals did not impair the chloroplast of *B. lambarenensis*. Samples irrigated with 5mg/l & 10mg/l of Cd , Fe and Va had lower chlorophyll a con tents that were statistically different at 0.05% and 0.01% levels of significance than those of the control. This shows that the chloroplasts were affected by these treatments. The worst effect was manifested in regime treated with Va.the effects of these heavy metals on chlorophyll a of *B. lambarenensis* was Va> Cd >Fe >Cu > Pb (Table 2)

Table 3 shows the effects of different heavy metals on the chlorophyll b contents of *B. lambarenensis*. Generally, chlorophyll b was generally less than chlorophyll a contents of this plant. Like chlorophyll a,, chlorophyll b contents of the treated and the control followed same trend of an increase from 1<sup>st</sup> week to the 5<sup>th</sup> week which subsequently reduced to the end of the experiment (Table 3). Samples irrigated with 5mg/l & 10mg/l of Cu & Cd and 10mgll of Fe had greater chlorophyll b than the control during the 1<sup>st</sup> week of experiment. This is also true for 5mg/l Cu, Fe & Pb and 10mgll of Fe

for the 2<sup>nd</sup> week of experiment. Statistical analysis revealed that the chlorophyll b contents of samples irrigated with 5mg/l of Cu, Pb and Va and 10mg/l of Pb were statistically the same with the control though less numerically. Moreover, the 5mg/l of Cd, Fe and 10mg/l of Fe were found to be statistically less than that of the control at 5% level of significance while those of 10mg/l of Cd & Va were statistically less than the control at both 5% and 1% levels of significance. The trend of toxicity on chlorophyll b contents was Cd>Fe>Va>Pb>Cu> control (Table 3).

The effects of different heavy metals on the total chlorophyll contents of *B. lambarenensis* are shown in Table 4. The total chlorophyll contents of samples irrigated with 5mg/l of Cd, Fe, Pb and 10mg/l of Fe were greater than the control during the 1<sup>st</sup> week while

those of 5mg/l & 10mg/l of Cu were greater from 2<sup>nd</sup> to the 4<sup>th</sup> week (Table 4). Samples of all the regimes showed same trend, an increase from 1<sup>st</sup> week to 5<sup>th</sup> week which subsequently reduced to the end of experiment. The total chlorophyll content of samples irrigated with 5mg/l & 10mg/l of Cu and Pb were found to be statistically the same with the control. Those irrigated with 5mg/l & 10mg/l of Cd, Va & Fe were statistically lower than the control (Table 4). The trend of action was Cd>Va>Fe>Pb>Cu> control. Generally, all the treatments had negative effects on the chloroplasts leading to lower contents.

## DISCUSSION

The changes in the green colouration and the distortion of the foliar cells and nuclei of *Barbula lambarenensis* as observed in this study is a manifestation of the detrimental effects of the heavy metals used. This fact is hinged on the maintenance of the bright green colouration, and intact cells of the control experiment. Sergio (1987) indicted pollutants such as SO<sub>2</sub> and heavy metals in the disappearance and changes in the flora around urban and industrial areas. Neolith Chemical Ltd (1988) directed that aqueous solutions of heavy metals such as lead, bitumeth, copper or mercury is necessary to prevent moss growth on masonry substrates. This further shows the effects of the heavy metals. Martin & Coughtrey (1982) reported that metals vary in their toxicity and degree of toxicity is proportional to their concentration. This is also observed in this study as the 10mg/l was found to be more toxic than 5mg/l of heavy metal solution used. The results observed in this study compared favourably with the findings of Gulvag *et al* (1974). Brown *et al* (1986) observed that low concentration of mercury inhibits photosynthesis temporarily, increases respiration, reduces chlorophylls a and b concentrations and also causes substantial loss of intracellular potassium from *Rhytidiadelphus squarrosus*. The reduction of chlorophylls a and b and total chlorophylls observed in this study agreed with the findings of Brown & Whitehead (1986). These reductions would certainly reduce the photosynthetic process of this plant.

Copper contaminated soils (10mg/g) was found to inhibit the protonemal growth of *Funaria hygrometrica* (Richardson, 1981). He further stated that SO<sub>2</sub> progressively changed green colouration of lichens and mosses to whitish brown as one moved from mildly to the highly polluted zone. A resultant colour change was observed in *Fontinalis squamosa* which absorbed metal ions from contaminated water (McLean and Jones, 1975). Farooqui *et al* (1993) reported that lead, cadmium, chromium, copper, zinc and aluminum are phytotoxic. This also confirmed the toxicity of Pb, Cd and Cu which were used in this study. The initial higher chlorophyll contents observed in some samples could be due to some damage done to the chloroplasts.

Mosses usually have efficient mechanism for absorbing metals and other pollutants from their environments with or without any destructive impact. This puts them at an advantage over other plants. The greater part of the metal contents of mosses is however not taken up and is not located within the cells but accumulates extracellularly (Richardson, 1981). The use of mosses as bio indicator could be attributed to their tolerance to heavy metals (Kakulu, 1993) and the ability to accumulate toxic elements (Brown, 1984; Rao, 1982;

Kovacs *et al.*, 1993). Brown (1984) reported that metals accumulate in form of particulate matters on the surfaces of the gametophores which are rough, providing many crevices for these tiny particles. Moreover, extracellular cation exchange capacity of bryophytes helps in explaining their metal uptake ((Brown and Bates, 1998).

## REFERENCES

- Annon, D I. (1949).Copper enzymes in isolated chloroplasts. Polyphenoloxidases in *Beta vulgaris* .*Plant Physiology* 24: 1-11.
- Bates, J.W. (2000). Mineral nutrition, substraction ecology and pollution. In Shaw, A.J., Goffinet, B.eds. *Bryophyte Biology*, Cambridge, New York, Cambridge Unipress, pg 248-311.
- Bates, J.W.(2002). Effects on bryophytes and lichens, In: Bell, J.N.B., Treshow, M. eds. *Air\_The\_Experimental Biology of Bryophytes: Pollution and Plant Life* 2<sup>nd</sup> edition. Chichester, John Wiley, pg 309-342
- Brown, D H. (1984). Uptake of mineral elements and their use in pollution monitoring. In:ed. A.S. Dyr. Academic Press New York, Pg. 367-374.
- Brown, D.H and Whitehead, A (1986). The effect of mercury on the physiology of *Rhytidiadelphus squarrosus* (Hedw). Warnst. *Journal of Bryology*, 14:367-374.
- Chakraborty, S.; Paratkar, G.T. and Kets ,V.G.(2006). Biomonitoring of trace element. Air pollution using mosses. *Aerosol and Air Quality Research* 6(3) :247-258.
- Gilbert, O.L. (1970) Further studies on the effects of sulphur dioxide on lichens and bryophytes. *New Phytol*,69; 605-627.
- Gradzinski, K (1990). Long-term ecological monitoring in the Parks of Poland. In: *Ecological Risks: Perspectives from Poland and United States* National Academic Press. pg. 252.
- Gulvag, B.M; Skaar, h and Ophus, E.M. (1974). An ultrastructural study of lead accumulation with leaves of *Rhytidiadelphus squarrossus* (Hedw)Warnst .*Journal of Bryology*,40; 311-317.
- Farooqui, P.A; Kulstresta, K; Mistra, J.; Pandey, V; Singh, N. and Sri Vastara, K. (1993). Heavy metals pollutants and plants. *Applied Botany Abstracts*: 13. 41-56.
- Fatoba, P.O. (1998).Reproductive phenology of three selected tropical African mosses in south western Nigeria. *Journal of Botany*, 11; 25-33.
- Kakulu, S.E. (1993). Biological monitoring of atmospheric trace metal deposition in North- Eastern Nigeria. *Environment Monitoring and Assessment*, 28:137-143.
- Kasanen, P. and VenetVaara, F.T.(1991).Comparison of biological collection of airborne heavy metals near ferrochrome and steel work. *Water Airr and Soil Pollution*, 60: 337-359.
- Kovacs,M.; Kaszab,I; Penksza,K; Szoke, P and Turosanyi, G.(1993) .Multi–elemental analysis of the species in a forest in the Hungarian forest steppe.*Phytocoenologia*, 23: 357.
- Rao, D.N. (1984). Response of bryophytes to air pollution. Smith A.J.E .Eds Chapman and Hall, London, pg 4 45-471.
- Martin, M. H. and Coughtrey, P.J.(1982 .*Biological monitoring of heavy metals pollution*.Applied Science Publishers, London and New York
- Mclean, R.O. and Jones, A.K.(1975).Studies of tolerance to heavy metals in the flora of the River Ystwyth and Clarachi, Wales. *Fresh Water Biology*, 5: 431-444.
- Mutkherjee, A. and Nuorteva, C.(1984). Relation between industrialization and growth of epiphytic lichens and mosses in Montreal. Canadaian Joutnal of Botany, 48:1480-14 ghtrey.

Neolith Chemicals Ltd (1988). Process for controlling growth of organisms on masonry. British Patent Application No 22205497, 7pp.

Richardson, D.H.S. (1981) *The biology of mosses*. Blackwell Scientific Publications. Oxford,. London, Edinburg, Boston, Melbourne. Pg 171-161.

Sergio,C.(1987).Epiphytic bryophytes and air quantity in the Jejo estuary. *Symposia. Biologica Hungarica*,35:795-801.

Strain H. H. and Svec, N. A. (1966). Extraction, separation and isolation of chlorophylls In *The chlorophyll eds* L. P. Varson and G.R. Seedy, Academic Press, New York. Pg 24-6.

**Table 2.** Effect of different heavy metals on chlorophyll a content of *Barbula lambarenensis*.

Concentration of Metal (mg/l)	Weekly Concentration of Chlorophyll (mg/g)												
	Week												× ± SD
	1	2	3	4	5	6	7	8	9	10	11	12	
Control	3.272	5.459	4.297	10.391	10.413	9.707	7.399	7.029	6.126	3.395	3.231	3.512	6.185± 1.293
Cu	2.894	10.461	5.973	15.764	9.354	8.527	4.374	3.219	3.113	2.396	2.304	2.862	5.936± 2.059 <sup>a</sup>
Cd	5.355	3.944	4.343	6.484	4.309	2.514	2.597	2.397	1.235	1.043	0.977	1.049	3.020± 1.361
5 Fe	5.330	4.443	6.541	6.667	7.834	6.088	4.356	2.822	1.534	0.690	1.379	1.489	4.097± 1.566
Pb	4.594	5.466	10.132	9.021	7.063	6.090	6.433	5.234	3.364	2.947	2.533	3.308	5.515± 1.548 <sup>a</sup>
Va	2.584	3.974	3.943	7.548	3.799	2.690	2.764	1.559	1.134	0.898	0.714	0.830	2.703± 1.401
Cu	2.811	11.127	6.314	11.069	6.823	4.995	5.447	3.974	3.645	3.357	3.025	3.229	5.484± 1.711 <sup>a</sup>
Cd	2.357	1.054	2.205	2.550	3.633	1.334	2.268	1.192	0.900	0.767	0.752	0.934	1.066± 0.958
10 Fe	6.766	7.243	2.890	4.086	7.147	3.571	2.810	2.214	1.725	1.722	1.396	1.577	3.595± 1.496
Pb	3.397	4.254	2.854	6.400	3.887	5.690	9.453	8.311	5.802	4.533	3.825	4.331	5.228± 1.146 <sup>a</sup>
Va	2.582	1.356	2.076	6.022	2.123	0.806	1.214	1.050	1.123	0.808	0.740	0.896	1.729± 1.224

Values with same superscript are statistically the same at 0.05 level of probability.

**Table 3.** Effect of different heavy metals on chlorophyll b content of *Barbula lambarenensis*.

Concentration of Metal (mg/l)	Weekly Concentration of chlorophyll (mg/g)												
	Week												
	1	2	3	4	5	6	7	8	9	10	11	12	× ± SD
Control	1.620	2.961	2.096	6.0731	5.4733	3.214	3.517	2.426	2.868	3.678	3.519	3.919	3.447± 1.133
Cu	1.737	5.587	3.144	8.124	5.185	3.013	2.068	2.275	1.832	1.833	1.692	1.596	3.173± 1.436 <sup>a</sup>
Cd	2.298	2.621	2.430	3.241	7.065	3.836	1.781	1.723	0.552	0.119	0.056	0.270	2.166± 1.409
5 Fe	1.585	3.029	3.066	3.436	4.671	2.882	2.732	3.275	0.486	0.148	0.148	0.349	2.150± 1.214
Pb	1.636	3.418	4.043	4.913	4.517	3.079	2.427	2.606	2.177	0.556	0.358	0.489	2.648± 1.249 <sup>a</sup>
Va	0.756	2.676	2.008	4.733	6.859	5.537	2.223	2.448	0.786	0.523	0.523	0.482	2.446± 1.476
Cu	1.921	5.911	3.654	5.889	4.803	1.949	4.105	3.815	1.964	1.773	1.570	1.570	3.233± 1.296 <sup>a</sup>
Cd	1.650	1.473	0.874	1.557	3.722	2.706	1.793	1.948	0.916	0.812	0.730	0.838	1.584± 0.947
10 Fe	1.559	4.845	0.731	3.686	4.420	3.525	2.447	2.251	1.029	0.573	0.624	1.565	2.271± 1.232
Pb	1.070	2.879	2.694	4.367	3.628	2.963	5.060	3.509	3.251	1.747	1.460	2.018	2.887± 1.088 <sup>a</sup>
Va	1.790	2.107	0.943	4.068	6.951	3.274	1.348	1.190	0.265	0.249	0.208	0.358	1.895± 1.418

Values with same superscript are statistically the same at 0.05 level of probability.

**Table 4.** Effect of different heavy metals on total chlorophyll content of *Barbula lambarenensis*.

Concentration of Metal (mg/l)	Weekly Concentration of Total chlorophyll (mg/g)												
	Week												
	1	2	3	4	5	6	7	8	9	10	11	12	× ± SD
Control	4.124	7.105	5.389	13.904	13.400	10.846	9.197	7.940	7.577	6.018	5.744	6.326	8.130± 1.770
Cu	3.912	13.539	7.690	20.145	12.270	9.694	5.427	4.650	4.177	3.584	3.384	3.763	7.686± 2.297 <sup>a</sup>
Cd	6.441	5.553	5.717	8.198	9.727	4.962	3.704	3.488	1.505	0.968	0.859	1.105	4.352± 1.715
5 Fe	5.798	6.322	8.093	8.520	10.563	7.558	5.991	5.193	1.695	1.323	1.273	1.539	5.322± 1.786
Pb	5.234	7.509	11.621	11.758	9.790	7.730	7.447	6.609	4.686	2.928	2.412	1.560	6.632± 1.854 <sup>a</sup>
Va	2.801	5.625	5.018	10.380	9.124	7.054	4.228	3.425	1.626	1.200	0.878	1.108	4.372± 1.785
Cu	4.004	14.373	8.417	14.306	9.840	5.800	8.091	6.618	4.733	4.014	3.769	4.045	7.334± 1.957 <sup>a</sup>
Cd	3.392	2.157	2.589	3.471	6.254	3.464	3.442	2.686	1.545	1.343	1.260	1.505	2.759±

10	Fe	6.967	1.025	3.033	6.599	9.775	6.031	4.462	3.796	2.327	2.075	1.701	2.671	1.186
	Pb	3.748	6.035	4.713	9.110	6.383	7.298	12.244	9.947	7.642	5.279	4.443	5.349	4.971± 1.715
	Va	3.692	2.960	2.543	8.536	7.816	3.520	2.181	1.967	1.162	0.888	0.796	1.033	6.849± 1.589 <sup>a</sup> 3.091± 1.605

Values with same superscript are statistically the same at 0.05 level of probability.